

# Discrete & Computational Geometry

## A Ten-Face Non-Edge-Sharing Wing Set on the Regular Icosahedron and a Decagonal Equatorial Balance --Manuscript Draft--

<b>Manuscript Number:</b>	DCGE-D-26-00035
<b>Full Title:</b>	A Ten-Face Non-Edge-Sharing Wing Set on the Regular Icosahedron and a Decagonal Equatorial Balance
<b>Article Type:</b>	Original Article
<b>Corresponding Author:</b>	YoungJune Jeon, B.S. Geowind Inc. Seoul, Seoul KOREA, REPUBLIC OF
<b>Corresponding Author Secondary Information:</b>	
<b>Corresponding Author's Institution:</b>	Geowind Inc.
<b>Corresponding Author's Secondary Institution:</b>	
<b>First Author:</b>	YoungJune Jeon, B.S.
<b>First Author Secondary Information:</b>	
<b>Order of Authors:</b>	YoungJune Jeon, B.S.
<b>Order of Authors Secondary Information:</b>	
<b>Funding Information:</b>	
<b>Abstract:</b>	We formalize a ten-face triangular wing set on a regular icosahedron with labeled vertices N, S, U1-U5, and L1-L5 and rotation axis NS. Each face is an isosceles 36-36-108 triangle anchored at a pole, and no two faces share an edge (vertex sharing is allowed). The ten pole-opposite intersection points on the equatorial plane form a regular decagon. The decagon radius equals $(\phi/2)$ times the icosahedron edge length, where $\phi$ is the golden ratio. We provide a reproducible construction and validation workflow suitable for CAD implementation.
<b>Suggested Reviewers:</b>	
<b>Author Comments:</b>	Data Availability: No datasets were generated or analyzed in this study. All constructions and validation steps are fully described in the manuscript and figures.

# A Ten-Face Non-Edge-Sharing Wing Set on the Regular Icosahedron and a Decagonal Equatorial Balance

YoungJune Jeon  
GeoWind

February 7, 2026

## Abstract

We formalize a ten-face triangular “wing” set on a regular icosahedron under a vertex labeling  $\{N, S, U_1, \dots, U_5, L_1, \dots, L_5\}$  with rotation axis  $NS$ . The wing faces satisfy: (i) each face is an isosceles  $36^\circ\text{--}36^\circ\text{--}108^\circ$  triangle with a  $36^\circ$  angle anchored at a pole ( $N$  or  $S$ ); (ii) distinct faces may share vertices but share no edges; and (iii) a natural equatorial cross-section yields a perfectly balanced regular decagon. We derive a closed form for the decagon radius,  $R = \frac{\varphi}{2}\ell$ , where  $\ell$  is the icosahedron edge length and  $\varphi = \frac{1+\sqrt{5}}{2}$  is the golden ratio. Beyond the geometric results, we interpret the “ten”-face closure as a symmetry-consistent design principle for a pole-anchored wing layout and provide a reproducible construction workflow.

## 1 Introduction

Regular polyhedra provide highly constrained geometric structures where design rules can be stated as axioms and verified as theorems. This paper defines and analyzes a specific triangular face set motivated by a GeoWind wing rule-set. Our first goal is mathematical: a precise definition and closed-form consequences for angles, symmetry, and a cross-sectional balance. Our second goal is design-facing: to express the same structure as a reproducible, checkable rule-set suitable for implementation in CAD/parametric geometry.

At a high level, the construction selects ten “golden” isosceles triangles (angles  $36^\circ\text{--}36^\circ\text{--}108^\circ$ ) whose small angle is anchored at one of two poles on the rotation axis, and whose edges are pairwise disjoint (edge non-sharing). When all ten faces are present, their pole-opposite edges induce a regular decagon on the equatorial plane with a radius determined by the golden ratio.

## 2 Preliminaries

Let  $\varphi = \frac{1+\sqrt{5}}{2}$  denote the golden ratio, satisfying  $\varphi^2 = \varphi + 1$ . Let  $I$  be a regular icosahedron with edge length  $\ell$  and center  $O$ . Choose opposite vertices  $N$  and  $S$  and call the line  $NS$  the rotation axis.

### 2.1 A standard coordinate model

For proofs that require explicit distance computations, we use the standard coordinate model of the icosahedron with edge length 2 whose vertices are

$$(0, \pm 1, \pm \varphi), \quad (\pm 1, \pm \varphi, 0), \quad (\pm \varphi, 0, \pm 1). \quad (1)$$

Scaling by  $\ell/2$  converts this model to edge length  $\ell$ .

### 3 Labeling and Wing-Face Axioms

**Definition 1** (Vertex labeling). *Label the 12 vertices of the icosahedron as*

$$V = \{N, S, U_1, \dots, U_5, L_1, \dots, L_5\},$$

where indices are taken modulo 5. The axis is the line  $NS$ .

**Definition 2** (GeoWind ten-face wing set). *Define the wing-face set  $\mathcal{F}$  as the following ten triangles:*

$$\begin{aligned} F_S(i) &= \triangle(S, U_i, L_i), & i &= 1, \dots, 5, \\ F_N(i) &= \triangle(N, U_i, L_{i-1}), & i &= 1, \dots, 5, \end{aligned}$$

with  $L_0 := L_5$ .

**Axiom 1** (Edge non-sharing; vertex sharing allowed). *For any two distinct faces  $F_a \neq F_b \in \mathcal{F}$ ,*

$$E(F_a) \cap E(F_b) = \emptyset,$$

*i.e., no edge is shared between two faces. Vertex sharing is allowed.*

**Axiom 2** (Non-intersection (geometric layout)). *The interiors of distinct faces do not intersect: for  $F_a \neq F_b \in \mathcal{F}$ ,*

$$\text{int}(F_a) \cap \text{int}(F_b) = \emptyset.$$

#### 3.1 Figures




Figure 1: Labeled regular icosahedron with axis  $NS$  and rings  $U_1, \dots, U_5, L_1, \dots, L_5$ .




Figure 2: Ten-face wing set  $\mathcal{F}$  embedded in the labeled icosahedron.




Figure 3: Equatorial plane  $\Pi$  and the regular decagon formed by the representative points  $p(F)$ .

## 4 Main Results

### 4.1 Angle structure via a golden-distance lemma

**Lemma 1** (Edge and diagonal distances in the standard model). *In the standard coordinate model (1) (edge length 2), (i) adjacent vertices have distance 2, and (ii) for an appropriate choice of opposite poles  $N, S$  and a corresponding upper vertex  $U$ , the distance between a pole and an adjacent upper vertex is 2, whereas the distance between the opposite pole and that upper vertex equals  $2\varphi$ . Equivalently, after scaling to edge length  $\ell$ , the corresponding distances are  $\ell$  and  $\varphi\ell$ .*

*Proof.* For (i), one may verify adjacency by direct computation for a representative pair. Let

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

$A = (0, 1, \varphi)$  and  $B = (1, \varphi, 0)$ ; then

$$\|A - B\|^2 = (0 - 1)^2 + (1 - \varphi)^2 + (\varphi - 0)^2 = 1 + (1 - 2\varphi + \varphi^2) + \varphi^2.$$

Using  $\varphi^2 = \varphi + 1$ , the expression becomes

$$1 + (1 - 2\varphi + \varphi + 1) + (\varphi + 1) = 4,$$

hence  $\|A - B\| = 2$ . Similar checks generate the full edge set by symmetry.

For (ii), take poles  $N = (0, 1, \varphi)$  and  $S = (0, -1, -\varphi)$  (these are opposite in (1)). Let  $U = (1, \varphi, 0)$ , which is adjacent to  $N$  by (i). Then

$$\|N - U\| = 2, \quad \|S - U\|^2 = (0 - 1)^2 + (-1 - \varphi)^2 + (-\varphi - 0)^2.$$

Compute

$$\|S - U\|^2 = 1 + (1 + 2\varphi + \varphi^2) + \varphi^2 = 2 + 2\varphi + 2\varphi^2.$$

Substituting  $\varphi^2 = \varphi + 1$  gives

$$\|S - U\|^2 = 2 + 2\varphi + 2(\varphi + 1) = 4 + 4\varphi = 4\varphi^2,$$

so  $\|S - U\| = 2\varphi$ . Scaling the coordinate model by factor  $\ell/2$  scales all distances by  $\ell/2$ , yielding  $\ell$  and  $\varphi\ell$ .  $\square$

**Lemma 2** (Cosine-law characterization of the golden gnomon). *Let a triangle have side lengths  $(\ell, \ell, \varphi\ell)$ . Then its angles are  $(36^\circ, 36^\circ, 108^\circ)$ .*

*Proof.* Let the equal sides have length  $\ell$ , and let the base have length  $\varphi\ell$ . By the law of cosines, the angle  $\theta$  opposite the base satisfies

$$\cos \theta = \frac{\ell^2 + \ell^2 - (\varphi\ell)^2}{2\ell \cdot \ell} = \frac{2 - \varphi^2}{2}.$$

Using  $\varphi^2 = \varphi + 1$  gives

$$\cos \theta = \frac{2 - (\varphi + 1)}{2} = \frac{1 - \varphi}{2} = -\cos 72^\circ,$$

hence  $\theta = 108^\circ$ . The remaining two angles are equal and sum to  $72^\circ$ , so each is  $36^\circ$ .  $\square$

**Theorem 1** (Golden-triangle shape:  $36^\circ-36^\circ-108^\circ$ ). *Every face  $F \in \mathcal{F}$  is an isosceles triangle with interior angles  $(36^\circ, 36^\circ, 108^\circ)$ . Moreover, for  $F_S(i)$  a  $36^\circ$  angle occurs at the pole  $S$ , and for  $F_N(i)$  a  $36^\circ$  angle occurs at  $N$ .*

*Proof.* Fix  $i$  and consider  $F_S(i) = \triangle(S, U_i, L_i)$ . In the regular icosahedron,  $S$  is adjacent to  $L_i$  and  $U_i$  is adjacent to  $L_i$ , so

$$|SL_i| = \ell, \quad |U_iL_i| = \ell.$$

By Lemma 1, the distance from  $S$  to the corresponding upper vertex is  $\varphi\ell$ , so

$$|SU_i| = \varphi\ell.$$

Thus  $F_S(i)$  has side lengths  $(\ell, \ell, \varphi\ell)$ , and by Lemma 2 its angles are  $(36^\circ, 36^\circ, 108^\circ)$  with a  $36^\circ$  angle at  $S$ .

For  $F_N(i) = \triangle(N, U_i, L_{i-1})$ ,  $N$  is adjacent to  $U_i$  and  $U_i$  is adjacent to  $L_{i-1}$ , hence

$$|NU_i| = \ell, \quad |U_iL_{i-1}| = \ell.$$

Again by Lemma 1, the remaining side has length  $\varphi\ell$ , so the same angle conclusion follows, with a  $36^\circ$  angle anchored at  $N$ .  $\square$

## 4.2 Edge non-sharing and maximality

**Theorem 2** (No shared edges within the ten faces). *The ten faces in  $\mathcal{F}$  satisfy Axiom 1: no two distinct faces share an edge.*

*Proof.* Each south face  $F_S(i)$  has edge set  $\{SU_i, SL_i, U_iL_i\}$ . For  $i \neq j$  these edges have different endpoints and thus cannot coincide. Each north face  $F_N(i)$  has edge set  $\{NU_i, NL_{i-1}, U_iL_{i-1}\}$ , and the same endpoint argument applies.

It remains to compare a south edge with a north edge. Any south edge incident to  $S$  cannot equal a north edge incident to  $N$ . A south edge of type  $U_iL_i$  cannot equal a north edge of type  $U_jL_{j-1}$  because equality would require  $U_i = U_j$  and  $L_i = L_{j-1}$ , hence  $i = j = j - 1 \pmod{5}$ , impossible. Therefore no edges are shared.  $\square$

**Theorem 3** (Maximality under pole-anchored edge non-sharing). *Consider pole-anchored wing triangles of the form  $\triangle(S, U_i, \cdot)$  whose edges must be pairwise non-shared. Then at most five such triangles can be anchored at  $S$ . Likewise, at most five can be anchored at  $N$ . Consequently, under pole anchoring at both poles and edge non-sharing, a construction can contain at most ten faces.*

*Proof.* Any triangle anchored at  $S$  using an upper vertex  $U_i$  necessarily contains the edge  $SU_i$ . Under edge non-sharing, no two such triangles may use the same edge  $SU_i$ . Since there are only five distinct upper vertices  $U_1, \dots, U_5$ , there are only five distinct edges  $SU_i$ , hence at most five edge-disjoint pole-anchored triangles can be anchored at  $S$ . The same argument holds for  $N$ .  $\square$

## 4.3 Equatorial decagon and closed-form radius

**Definition 3** (Equatorial plane and cross-edge midpoints). *Let the equatorial plane be*

$$\Pi := \{x \in \mathbb{R}^3 : (x - O) \cdot (N - S) = 0\},$$

*i.e., the plane through  $O$  perpendicular to axis  $NS$ . For a face  $F$ , let  $c(F)$  denote the edge not incident to the corresponding pole (the “cross-edge”). Define the representative point*

$$p(F) := \text{mid}(c(F)),$$

*the midpoint of the cross-edge.*

**Proposition 1** (Cross-edges for the ten faces). *For the faces in Definition 2,*

$$c(F_S(i)) = U_iL_i, \quad c(F_N(i)) = U_iL_{i-1}.$$

*Proof.* In  $F_S(i) = \triangle(S, U_i, L_i)$  the two edges incident to pole  $S$  are  $SU_i$  and  $SL_i$ , hence the remaining edge is  $U_iL_i$ . The north case is identical.  $\square$

**Theorem 4** (Perfect circular balance: regular decagon on  $\Pi$ ). *The set  $\{p(F) \mid F \in \mathcal{F}\} \subset \Pi$  consists of ten points lying on a single circle centered at  $O$ , forming a regular decagon with  $36^\circ$  angular spacing.*

*Proof.* By Proposition 1, each  $p(F)$  is the midpoint of a  $U$ - $L$  edge. By symmetry about the axis and the central plane through  $O$  orthogonal to  $NS$ , each such midpoint lies in  $\Pi$ , so  $p(F) \in \Pi$ .

Next, consider the two midpoint sets

$$P_1 = \{\text{mid}(U_iL_i)\}_{i=1}^5, \quad P_2 = \{\text{mid}(U_iL_{i-1})\}_{i=1}^5.$$

Rotation by  $72^\circ$  about axis  $NS$  maps  $U_i \mapsto U_{i+1}$  and  $L_i \mapsto L_{i+1}$ , so it cyclically permutes  $P_1$  and also  $P_2$ . Hence each of  $P_1$  and  $P_2$  forms a regular pentagon on a circle centered at  $O$  in  $\Pi$ .

To show that  $P_1 \cup P_2$  is a regular decagon, it suffices to show a  $36^\circ$  phase offset between one point in  $P_1$  and a neighboring point in  $P_2$ . This is established in Proposition 2. Therefore the two pentagons interlace into a regular decagon with  $36^\circ$  spacing.  $\square$

**Proposition 2** (Phase offset:  $\cos 36^\circ = \varphi/2$ ). In the standard model (1) (edge length 2), let

$$M = \text{mid}((0, 1, \varphi), (1, \varphi, 0)), \quad M' = \text{mid}((0, 1, \varphi), (-1, \varphi, 0)).$$

Then  $\angle MOM' = 36^\circ$  and

$$\frac{M \cdot M'}{\|M\| \|M'\|} = \frac{\varphi}{2}.$$

*Proof.* Compute

$$M = \left(\frac{1}{2}, \frac{1+\varphi}{2}, \frac{\varphi}{2}\right), \quad M' = \left(-\frac{1}{2}, \frac{1+\varphi}{2}, \frac{\varphi}{2}\right).$$

Their dot product is

$$M \cdot M' = -\frac{1}{4} + \left(\frac{1+\varphi}{2}\right)^2 + \left(\frac{\varphi}{2}\right)^2.$$

Using  $\varphi^2 = \varphi + 1$ ,

$$\left(\frac{1+\varphi}{2}\right)^2 = \frac{1+2\varphi+\varphi^2}{4} = \frac{2+3\varphi}{4}, \quad \left(\frac{\varphi}{2}\right)^2 = \frac{\varphi^2}{4} = \frac{\varphi+1}{4}.$$

Therefore

$$M \cdot M' = -\frac{1}{4} + \frac{2+3\varphi}{4} + \frac{\varphi+1}{4} = \frac{2+4\varphi}{4} = \frac{1+2\varphi}{2}.$$

Meanwhile,  $\|M\| = \|M'\| = \varphi$  (see Theorem 5), so  $\|M\| \|M'\| = \varphi^2$ . Thus

$$\frac{M \cdot M'}{\|M\| \|M'\|} = \frac{(1+2\varphi)/2}{\varphi^2} = \frac{\varphi}{2},$$

which equals  $\cos 36^\circ$ . Hence  $\angle MOM' = 36^\circ$ .  $\square$

**Theorem 5** (Closed-form radius  $R = \frac{\varphi}{2}\ell$ ). Let  $R$  be the radius of the circle in Theorem 4. Then

$$R = \frac{\varphi}{2}\ell.$$

*Proof.* In the standard model (1) (edge length 2), consider adjacent vertices  $A = (0, 1, \varphi)$  and  $B = (1, \varphi, 0)$ . Their midpoint is

$$M = \left(\frac{1}{2}, \frac{1+\varphi}{2}, \frac{\varphi}{2}\right).$$

Using  $\varphi^2 = \varphi + 1$ ,

$$\|M\|^2 = \left(\frac{1}{2}\right)^2 + \left(\frac{1+\varphi}{2}\right)^2 + \left(\frac{\varphi}{2}\right)^2 = \varphi^2,$$

so  $\|M\| = \varphi$ . Hence the midpoint circle in the edge-length-2 model has radius  $\varphi$ . Scaling to edge length  $\ell$  multiplies distances by  $\ell/2$ , giving  $R = \varphi(\ell/2) = \frac{\varphi}{2}\ell$ .  $\square$

**Corollary 1.** If  $\ell = 1$  m, then

$$R = \frac{\varphi}{2} \text{ m} \approx 0.809016994 \text{ m}.$$

## 5 Design Interpretation: Why “Ten”?

The number ten is not merely a count; it encodes a closure condition consistent with the icosahedron’s fivefold rotational symmetry about the pole axis. On each pole, there are exactly five distinct edges  $SU_i$  (and  $NU_i$ ). Under edge non-sharing, these edges act as five “ports” for pole-anchored faces (Theorem 3). The chosen index shift between south and north faces prevents edge duplication while interlacing the two pentagonal midpoint sets into a decagon (Theorem 4).

From a design standpoint, the equatorial decagon provides a compact diagnostic: if a CAD model implements the ten faces correctly, the midpoints of the pole-opposite edges must lie on a single circle and fall into a regular decagon with radius  $\frac{\varphi}{2}\ell$ . This offers a simple geometric “sanity check” independent of aerodynamic modeling.

## 6 Reproducible Construction Workflow (CAD/Parametric)

We summarize a practical procedure to generate and validate  $\mathcal{F}$ .

### Algorithm 1 (Wing-face generation and validation)

1. **Create a labeled icosahedron.** Fix edge length  $\ell$  and label vertices as in Definition 1.
2. **Generate faces.** Create  $F_S(i) = \triangle(S, U_i, L_i)$  and  $F_N(i) = \triangle(N, U_i, L_{i-1})$  for  $i = 1, \dots, 5$ .
3. **Edge non-sharing check.** List all edges of the ten triangles (unordered vertex pairs) and verify no duplicates.
4. **Angle check.** Verify side lengths  $(\ell, \ell, \varphi\ell)$  and angles  $(36^\circ, 36^\circ, 108^\circ)$  for each face.
5. **Equatorial decagon check.** Compute  $p(F)$  for all faces. Confirm decagon geometry and radius  $R = \frac{\varphi}{2}\ell$ .
6. **Non-intersection check (optional).** Perform triangle–triangle intersection tests to numerically verify Axiom 2.

## 7 Related Geometric Context

The appearance of  $\varphi$  in the icosahedron is classical: coordinate realizations, vertex/edge relations, and duality with the dodecahedron all involve the golden ratio. The present contribution is a specific edge-disjoint, pole-anchored ten-face rule-set whose equatorial midpoints close into a decagon with a simple closed-form radius. For general background, see Coxeter and Grünbaum.

## 8 Conclusion

We defined a pole-anchored ten-face wing set on a regular icosahedron and proved: (i) each face is a  $36^\circ - 36^\circ - 108^\circ$  isosceles (golden) triangle; (ii) faces share no edges and at most ten such pole-anchored edge-disjoint faces can exist across both poles; and (iii) the pole-opposite edge midpoints form a regular decagon on the equatorial plane with radius  $R = \frac{\varphi}{2}\ell$ . These results provide a mathematical foundation and a practical validation workflow for geometry-driven design implementations.

## References

- [1] H. S. M. Coxeter, *Regular Polytopes*, Dover Publications.
- [2] B. Grünbaum, *Convex Polytopes*, Springer.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65